





Know Diabetes by Heart™



Approximately 1 in 3 American adults with diabetes has CKD.



CVD risk is high in people with CKD.



Kidney disease is ranked in the top leading causes of death in the United States.



Every 24 hours, 170 people with diabetes begin treatment for kidney failure.

Screening for Diabetes-Related Kidney Disease



Who?

- ✓ Everyone with type 2 diabetes
- ✓ Everyone with type 1 diabetes for ≥ 5 years



How?

UACR and eGFR



When?

Annually


Management of Diabetes-Related Kidney Disease

Why Manage?

Decreases risk of CKD progression

Cardiovascular risk reduction

Pillars of Therapy to Reduce Cardiorenal Risk



CKD = Chronic kidney disease
 CVD = Cardiovascular disease
 UACR = Urinary albumin to creatinine ratio
 eGFR = Estimated glomerular filtration rate

RAAS Blockade = Renin-angiotensin-aldosterone system blockade
 SGLT2-inhibitors = Sodium-glucose cotransporter 2 inhibitor
 NS-MRAs = Nonsteroidal mineralocorticoid receptor antagonists
 GLP-1 RAs = Glucagon-like peptide 1 agonists

Learn more at [KnowDiabetesbyHeart.org](https://www.knowdiabetesbyheart.org)